

Q^2 -dependence of semi-inclusive electron-nucleus scattering and nucleon-nucleon correlations

O. Benhar ^a, S. Fantoni ^b, G.I. Lykasov ^c and N.V. Slavin^c

^a *INFN, Sezione Sanita', I-00161, Roma, Italy*

^b *Interdisciplinary Laboratory, SISSA,
and INFN, Sezione di Trieste, I-34014, Trieste, Italy*

^c *Joint Institute for Nuclear Research,
141980 Dubna, Moscow Region, Russia*

(February 1, 2008)

Abstract

We analyze semi-inclusive electron-nucleus processes $e + A \rightarrow e' + h + X$ at moderate Q^2 and energy transfer ν . Our results show that nucleons bound in the nuclear medium are distributed according to a function f_A that reduces to the standard light-cone distribution in the Bjorken limit and exhibits a sizeable Q^2 -dependence at lower Q^2 , particularly when $Q^2 \sim \nu^2$. This distribution function is employed to calculate the cross section of the reaction $e + A \rightarrow e' + p + X$, in the kinematical region in which the detected proton is emitted backward in the target rest frame.

PACS numbers: 13.60.Le, 25.30Fj, 25.30Rw

The semi-inclusive electron-nucleus scattering process $e + A \rightarrow e' + h + X$, in the kinematical region in which the detected hadron h is emitted approximatively backward in the target rest frame, has been recently investigated both experimentally [1] and theoretically [2]. These studies were mainly aimed at pinning down the role of the high-momentum tails of the nucleon momentum distribution in nuclear matter. In ref. [2] two approaches, based on different descriptions of nuclear structure at short interparticle distance, were compared. The first one was based on nuclear many-body theory, in which the nuclear target is described in terms of nonrelativistic structureless nucleons whose distribution in momentum and removal energy is dictated by the nuclear spectral function $P(\mathbf{k}, E)$. Within the alternative approach, on the other hand, the relativistic invariant phase space volume available to a quark in a nucleus was computed taking into account nucleon-nucleon correlations and the true Regge asymptotic of the quark distribution in the nucleon.

In the first approach discussed in ref. [2] the spectral function is used to generate the distribution of nucleons in the nuclear target, f_A , as a function of the relativistic invariant quantity

$$z = \frac{(kq)}{(P_A q)} \frac{M_A}{m} . \quad (1)$$

In the above equation $P_A \equiv (M_A, \mathbf{0})$, M_A being the target mass, and $k \equiv (k_0, \mathbf{k})$ denote the initial four-momenta of the target nucleus and the struck nucleon in the laboratory frame, respectively, $q \equiv (\nu, \mathbf{q})$ is the four-momentum transfer and m is the nucleon mass. The regime of deep inelastic scattering (DIS) is defined by the Bjorken limit $Q^2, \nu \rightarrow \infty$ with $x = Q^2/2m\nu$ finite ($Q^2 = |\mathbf{q}|^2 - \nu^2 = -q^2$). In this limit the variable z reduces to the ordinary light-cone variable:

$$z \rightarrow z_{Bj} = \frac{k^+}{m} = \frac{(k_0 - k_z)}{m} . \quad (2)$$

However, it has been recently pointed out [3] that when the value of Q^2 is not large enough and the Bjorken limit is not yet reached the quantity z cannot be identified with the light cone variable. As a consequence, the distribution function f_A depends upon both z and

$\beta = (|\mathbf{q}|/\nu) = (1 + 4m^2x^2/Q^2)^{1/2}$. The results of ref. [3] show that this β -dependence is sizeable, particularly at $z > 1$.

The semi-inclusive spectra of π mesons and protons discussed in ref. [1] have been produced scattering 14.5 GeV electrons and positrons off CO at moderate Q^2 and ν . The kinematical domain covered by the experiment corresponds to energy loss $0.2 < \nu < 12.5$ GeV and Q^2 in the range $0.1 < Q^2 < 5.0(\text{GeV}/c)^2$. Hence, the beta dependence of the distribution function f_A is expected to be relevant to the theoretical analysis of the data of ref. [1] and has to be quantitatively taken into account. In this note we extend the calculations of ref. [2], including the dependence of $f_A(z, \beta)$ upon β in the analysis of semi-inclusive $e + A \rightarrow e' + h + X$ processes at moderate Q^2 and ν .

The distribution function $f_A(z, \beta)$ is defined as:

$$f_A(z, \beta) = z \int d^4k S(k) \delta \left(z - \frac{(kq)}{(P_A q)} \frac{M_A}{m} \right), \quad (3)$$

where $S(k)$ denotes the relativistic vertex function, that can be approximated by the non-relativistic spectral function $P(\mathbf{k}, E)$ according to [4]

$$S(k) = \left(\frac{m}{k_0} \right) P(\mathbf{k}, E), \quad (4)$$

with

$$k_0 = M_A - [(M_A - m + E)^2 + |\mathbf{k}^2|]^{1/2}. \quad (5)$$

The details of the calculation of $f_A(z, \beta)$ are presented elsewhere [3] and will be omitted here for the sake of conciseness. The numerical results of ref. [3], corresponding to infinite nuclear matter at equilibrium density, are summarized in Fig. 1, where the z behaviour of the function $f_A(z, \beta)$ is shown for different values of β . It clearly appears that $f_A(z, \beta)$ strongly depends upon β at $z > 1$.

Let us now turn to the semi-inclusive momentum spectrum of the protons emitted in the $e + A \rightarrow e' + p + X$ around the backward direction, in the kinematical region forbidden to the quasi-free electron-nucleon process. At small transverse momenta, i.e. at $p_t \sim 0$,

this spectrum, that will be denoted $\rho(z, x, Q^2)$, is proportional to the distribution of the nucleons in the target as a function of the variable z , defined as in eq.(1). According to the two approaches discussed in ref. [2] this distribution can be identified with either the function $f_A(z, \beta)$ of eq.(3) or with a function $T_A(z)$, constructed under the assumption that the nucleus can be viewed as a collection of multiquark bags whose structure functions exhibit Regge asymptotic behaviour [5]. Within the framework of the latter approach the hadron yields measured in hadron-nucleus collisions in the kinematical region forbidden to hadron-nucleon inelastic scattering, the so-called limiting fragmentation processes, can be quantitatively accounted for [5–7]. However, in this picture the argument of the distribution function T_A is the usual light-cone variable z_{Bj} , that coincides with z of eq.(1) in the Bjorken limit only. One has:

$$\rho(z, x, Q^2) \equiv \int d^2p_t E_p \frac{d\sigma(e + A \rightarrow e' + p + X)}{d^3p dE_{e'} d\Omega_{e'}} \sim T_A(z_{Bj}) F_2^N(x, Q^2) , \quad (6)$$

where $F_2^N(x, Q^2)$ is the nucleon structure function. On the other side, the first approach discussed in ref. [2], based on the use of the nuclear spectral function, leads to the following expression:

$$\rho(z, x, Q^2) \sim f_A(z, Q^2) F_2^N(x, Q^2) , \quad (7)$$

with f_A defined as in eq.(3) (notice however that in the above equation, the dependence upon β has been replaced with an explicit dependence upon Q^2).

In order to compare the spectra defined by eqs.(6) and (7) to the data of ref. [1] one has to integrate over the relevant ranges of Q^2 and ν , as done in ref. [2]. The integrated spectra are shown in Fig. 2 as a function of the kinetic energy of the detected proton, T , emitted at angle $\Theta = \arccos(-0.55)$ with respect to the direction of the three-momentum transfer $\mathbf{q}$. The calculations have been performed using the empirical nucleon structure function $F_2^N(x, Q^2)$ at $0.2 < Q^2 < 5 \text{ (GeV/c)}^2$, extracted from electron-proton and electron-deuteron data according to the parametrization of ref. [8].

The solid line in Fig. 2 represents the integrated proton spectrum computed using eq.(7), with $f_A(z, \beta)$ given by eqs.(3)-(5), and the nuclear matter spectral function of ref. [9]. The

calculation has been carried out for proton kinetic energies $T \leq T_{max}$, with $T_{max} \sim 0.28$ GeV, that corresponds to $z \leq 1.9$. This cutoff reflects the momentum range $k \leq 0.8$ (GeV/c) covered by the spectral function of ref. [9]. The effect of the dependence of the distribution function upon β (or, equivalently, upon Q^2) can be estimated comparing the solid line to the dashed line, which has been obtained setting $\beta = 1$, i.e. identifying z of eq.(1) with the light-cone fraction of eq.(2). The two curves significantly differ, indicating that, as expected, in the moderate Q^2 range of the data of ref.[1] the Bjorken limit is not applicable and the Q^2 dependence associated with the distribution function cannot be disregarded.

In Fig. 2 we also show, for comparison, the integrated proton spectrum computed from eq.(6) with the distribution function T_A discussed in ref. [2]. Comparison between the solid and dot-dash lines suggests the existence of a connection between the processes $h+A \rightarrow p+X$ and $e+A \rightarrow e'+p+X$ when the produced proton is emitted backward, i.e. in the kinematical domain forbidden to hadron-nucleon and electron-nucleon collisions.

In conclusion, the results of our work, summarized in Figs. 1 and 2, show significant dependence of the nucleon distribution f_A upon the magnitude of the virtual photon velocity $\beta = |\mathbf{q}|/\nu$, or equivalently upon Q^2 , at fixed x . The inclusion of this effect in the theoretical analysis of semi-inclusive backward proton production in the kinematical regime of the experiment reported in ref.[1] leads to an improvement of the agreement between theory and data up to $T \sim 2.8$ GeV, corresponding to the maximum quantity $z(Q^2) \sim 1.9$. Unfortunately, the limitations inherent to the nonrelativistic nature of the many-body theory employed in the calculation of the spectral function make it impossible to extend the calculation of $f_A(z, \beta)$ to larger values of z .

The results obtained from eq.(6) using the nucleon distribution T_A dictated by true Regge asymptotic at $z_{Bj} \rightarrow 2$ and $z_{Bj} \rightarrow 3$ [5,6] are also in satisfactory agreement with the data at $T < 2.8$ GeV. In addition, they account for the qualitative behaviour of the experimental spectrum up to $T \sim 0.4$ GeV, corresponding to $z_{Bj} \sim 2.5$. A detailed theoretical analysis of the relationship between the functions f_A and T_A may therefore provide some guidance on the the extrapolation of the nuclear spectral function to the large k domain, inaccessible

within nonrelativistic nuclear many-body theory.

ACKNOWLEDGMENTS

This work had been encouraged and supported by the Russian Foundation of Fundamental Research. We gratefully acknowledge very helpful discussions with A. Fabrocini. One of us (GIL) wishes to thank S. Fantoni for the kind hospitality at the Interdisciplinary Laboratory of SISSA, where part of this work has been carried out.

REFERENCES

- [1] P.V. Degtyarenko *et al*, Phys.Rev C, **50**, R541 (1994).
- [2] O. Benhar, S. Fantoni, G.I. Lykasov and N.V. Slavin, Phys. Rev. C, **55**, 244 (1997).
- [3] O. Benhar, S. Fantoni, G.I. Lykasov and N.V. Slavin, preprint SISSA-ISAS 78/97/CM (1997).
- [4] O. Benhar, V.R. Pandharipande and I. Sick, Phys. Lett. B, in press.
- [5] A.V. Efremov, A.B. Kaidalov, V.T. Kim, G.I. Lykasov and N.V. Slavin, Sov. Journ. Nucl. Phys., **47**, 868 (1988).
- [6] G.I. Lykasov, Phys. Part. Nuclei, **24**, 59 (1993).
- [7] A.V. Efremov, A.B. Kaidalov, G.I. Lykasov and N.V. Slavin, Sov. Journ. Nucl. Phys., **57**, 874 (1994).
- [8] A. Capella, A. Kaidalov, C. Merion and J. Tran Than Van, Phys. Lett., **B337**, 358 (1994); **B343**, 403 (1995).
- [9] O. Benhar, A. Fabrocini and S. Fantoni, Nucl. Phys., **A505**, 267 (1989).

FIGURES

FIG. 1. z -dependence of the distribution function $f_A(z, \beta)$ of eq.(3) at different values of β . Solid line: $\beta = 1$; dashed line: $\beta = 1.1$; dot-dash line: $\beta = 1.2$; dotted line: $\beta = 1.3$.

FIG. 2. Kinetic energy spectra of protons emitted at angle Θ in the semi-inclusive reaction $e + CO \rightarrow e' + p + X$. The solid curve corresponds to the full calculation carried out using eq.(7), whereas the dashed line represents the results obtained setting $\beta=1$. The dot-dash line shows the spectrum calculated from eq.(6) using the nucleon distribution T_A of ref.[5]. The experimental data are taken from ref.[1].

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9707252v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9707252v1>